

International Journal of Environmental Sciences

(A peer reviewed International Journal)

Thermal Effect Due to Induced Field of Broadcasting Radiation

International Journal of Environmental Sciences, Vol. 1 No. 2, pp.50-55, 2277-1948, 2012

ISSN 2277 – 1948

Article type *Full Length Research Article*

Submission date *12 March 2012*

Acceptance date *30 March 2012*

Publication date *15 April 2012*

Article URL <http://www.crdeep.org/category/ijes>

Authors **Hemendra Tripathi and P.P. Pathak*

This peer-reviewed article was published immediately upon acceptance. It can be downloaded, printed and distributed freely for any purposes from CRDEEP website.

Hard copy of Journal is also available on request.

For information about publishing your research in CRDEEP International Journals please visit our website www.crdeep.org

© 2012. All Rights Reserved by CRDEEP

CRDEEP Head Office:315/10, Kaulagarh Road, Rajendranagar, Indervihar, Dehradun, 248001, Uttrakhand, India.



Full Length Research Paper**Thermal Effect Due to Induced Field of Broadcasting Radiation*****Hemendra Tripathi and P.P. Pathak****Department of Physics, Gurukula Kangri University, Haridwar-249404, India***Corresponding Author: Hemendra Tripathi***ABSTRACT**

Thermal effect inside the tissues causes due to penetration and absorption of radio frequency radiation emitted by radio broadcasting antenna. Numerical calculation is made for the specific absorption rate in consequence of induced electric field due to radio tower at different distances from the transmission tower. The depth of penetration inside the blood and cerebro spinal fluid is also varied. The temperature of the body is maintained at a set point through the thermoregulatory mechanism of the body, is also discussed. The damage of tissues may cause due to unavailability of sufficient mechanisms to carry away all of the unwanted heat during the gain of heat energy deposited through broadcasting antenna, which would be harmful for human being. The comparison is made by the international guidelines of World Health Organization (WHO) and International Commission on Non-Ionization Radiation Protection (ICNIRP) for the amount of energy deposition due to radio frequency of high power broadcasting antenna inside the selected tissues. To follow the international guidelines carefully for installation of high power transmitters of radio frequency radiation is recommended and the suggestion is made for the minimum distance of human being from broadcasting antenna for safe exposure

Keywords: *Induced electric field, Specific absorption rate, Rate of Change of temperature, Thermoregulatory mechanism, Broadcasting antenna.*

INTRODUCTION

In present times of faster communication, we are exposed to many types of radiation from all types of communication systems like TV, radio and mobile telephony etc. Generally speaking, some radiation penetrates and is being absorbed by the human body when it is exposed to it. Depending on the frequency of radiation, the human body interacts with such a field via induced currents and thermal effects. The field generated within the body, the so-called internal field, is determined by the amount by which a body is influenced by electromagnetic fields. The EMF can be characterized by several different parameters (field strength, field direction, field orientation, field complexity and so on) (Gandhi and Mohammad, 2008). The quantity used to measure how much RF energy is actually absorbed in a body is called the specific absorption rate (SAR). This SAR also varies with the dimension of tissues (Dein and Amr, 2010). The absorbed microwave energy produces molecular vibration and converts the energy into heat. When the rate of energy absorption is high, it produces heating in living tissues (Ozen *et al.* 2008). The heat generated in the medium is proportional to the absorbed power. The biological effects of radiofrequency energy depend on the rate at which power is absorbed (Osepchuk and Petersen, 2003). If the organism cannot dissipate this heat energy as fast as heat is produced, the internal temperature of the body will rise.

Exposure from TV and Radio transmitters have been studied by Joseph and Martens (2006) and Sirav and Seyhan (2009).

Pathak *et al.* (2003) and Kumar *et al.* (2008) assessed the temperature change in tissues per second by taking the TV transmitters of high frequency radiation as a point source. Due to exposure to microwave radiation many health risks have been studied including brain tumors, acoustic neuroma, leukemia and testicular cancer, sleep disturbances and headache (Johansen, 2004; Takebayashi *et al.* 2006, 2008; Hardell *et al.* 2007). Wessapan (2011) calculated the local SARs and temperature increase of human model for various operating frequencies, also focusing on the interaction between electromagnetic field and organs in the human trunk.

In this manuscript the specific absorption rate as a result of induced electric field is assessed due to the radio frequency antenna of All India Radio, situated in Mussoorie (Uttarakhand), India. The calculation of induced electric field and SAR in selected tissues is made by taking the transmitter of finite length dipole antenna. The frequency used for the calculation is 102.1 MHz of FM (rel.) broadcasting and power of the antenna is 10 kW. The calculation is made at different distances from the antenna and at different depth inside the tissues.

Public access to broadcasting antennas is normally restricted so individuals cannot be exposed to high-level fields that might exist near antennas. Ambient RF radiation levels in inhabited areas near broadcasting facilities must be typically well below the exposure levels recommended by current

standards and guidelines. Therefore, precautions must be taken to ensure that maintenance personnel are not exposed to unsafe RF fields.

Thermoregulation

Thermoregulation of the human body in the presence of RF fields follows the laws of thermodynamics in which heat and work are balanced and conserved. The whole gain energy becomes equal to the spend energy by the body. Thus there is no excess temperature in the body in this natural process. More specialized adaptation of the relevant equations has been developed by thermal physiologists. The balance of heat energy is expressed by the following equation;

$$M + W = E \pm R \pm C \pm D \pm S \quad , \quad (1) \text{ where}$$

M is the rate at which thermal energy is produced through metabolic processes, W , the rate at which the work is produced, E , the rate of exchange with the surroundings via evaporation, R , the rate of heat exchange with the surroundings via radiation, C , the rate of heat exchange with the environment via convection, D , the rate of heat exchange with the surroundings via conduction and S represents the rate of body heat storage.

The ability of various thermal sensors to initiate mechanisms of heat loss, during the gain of heat energy deposited by the dissipation of electrical currents deposited by RF energy depends on the availability of sufficient mechanisms (vasomotor, sudomotor in addition to convection and radiation) to carry away all of the unwanted heat. Without this, cooling cannot occur and the core body temperature will continue to rise.

Thus, when electromagnetic radiation is penetrated inside the body, the energy is absorbed by the tissues of the biological material. It works as a source of production of extra energy inside the body.

$$M \pm W + E_R = E \pm R \pm C \pm D \pm S \quad , (2) \text{ where}$$

E_R is the energy due to electromagnetic radiation of broadcasting antenna. Above equation becomes un-equilibrium because production of energy becomes greater to the energy inside the body. This excess energy may increase the temperature of the tissue and may harmful in much other way for tissue life.

MATERIAL AND METHODS

As we want to access the effect of broadcasting antenna, whose power P is the only parameter we know, we have to relate the electric field (more generally E_{rms}) to the power of

$$E_{rms} = \frac{\sqrt{90P}}{r} = 9.487 \frac{\sqrt{P}}{r} \quad , \quad (3)$$

transmitters which is distance far from the biological body (Prasad, 1999).

When a human body is exposed to the EM wave of electric field E_{rms} , it penetrates into the body. It results into inside or induced field E_i at a given depth z given by Polk (1996).

$$E_i = E_{rms} \exp\left(\frac{-z}{\delta}\right) \quad , \quad (4) \text{ where}$$

δ is skin depth, which is the distance over which the field decreases to 0.368 of its value just inside the boundary at angular frequency ω , given as

$$\delta = \frac{1}{\omega \sqrt{\frac{\mu\epsilon}{2} \left\{ \left[(1+p^2)^{\frac{1}{2}} - 1 \right] \right\}}} \quad , \quad (5) \text{ where}$$

μ is the permeability of body material, ϵ , its permittivity and biological material ratio $p = \frac{\sigma}{\epsilon\omega}$, σ being its conductivity.

It is generally accepted that the SAR is the most appropriate metric for determining electromagnetic exposure, i.e. the mass averaged rate of energy absorption in tissue, is related to the induced electric field E_i (V/m) can be determined at any point from the relation (Hirata et al., 2008)

$$SAR = \frac{\sigma E_i^2}{\rho} \quad , \quad (6) \text{ where}$$

σ is the conductivity of the tissues for which the calculation is made and ρ is their mass density.

CALCULATION

This electromagnetic wave penetrates inside the body. Thus, the induced electric field at different distances in the blood and cerebro spinal fluid of the human body at different depth is numerically evaluated in Table 1 and 2. The induction of field causes the deposition of energy in the form of SAR. The values of SAR corresponding to induced electric field are also calculated in Table 3 and 4. The conductivity of the body tissues is taken from Gabriel et al. (1996 a, b, c)

Table 1: Induced Electric Field (V/m) at different depths in blood at different distances due to radiation from All India Radio Antenna of power 10 kW at 102.1 MHz

Distance (m)	10 μm	20 μm	30 μm	40 μm	50 μm
10	94.8522	94.8344	94.8166	94.7988	94.7810
20	47.4261	47.4172	47.4083	47.3994	47.3905
30	31.6171	31.6111	31.6052	31.5993	31.5933
40	23.7135	23.7091	23.7046	23.7002	23.6957
50	18.9704	18.9669	18.9633	18.9598	18.9562
60	15.8090	15.8061	15.8031	15.8001	15.7972
70	13.5505	13.5479	13.5454	13.5428	13.5403
80	11.8568	11.8545	11.8523	11.8501	11.8479
90	10.5390	10.5370	10.5351	10.5331	10.5311
100	9.4852	9.4834	9.4817	9.4799	9.4781
110	8.6234	8.6218	8.6201	8.6185	8.6169
120	7.9045	7.9030	7.9015	7.9001	7.8986

Table 2: Induced Electric Field (V/m) at different depths in cerebro spinal fluid at different distances due to radiation from All India Radio Antenna of power 10 kW at 102.1 MHz

Distance (m)	10 μm	20 μm	30 μm	40 μm	50 μm
10	94.8454	94.8207	94.7961	94.7715	94.7469
20	47.4227	47.4104	47.3981	47.3858	47.3735
30	31.6148	31.6066	31.5984	31.5902	31.5820
40	23.7118	23.7057	23.6995	23.6934	23.6872
50	18.9691	18.9641	18.9592	18.9543	18.9494
60	15.8079	15.8038	15.7997	15.7956	15.7915
70	13.5495	13.5460	13.5424	13.5389	13.5354
80	11.8559	11.8528	11.8498	11.8467	11.8436
90	10.5383	10.5355	10.5328	10.5301	10.5273
100	9.4845	9.4821	9.4796	9.4772	9.4747
110	8.6228	8.6205	8.6183	8.6160	8.6138
120	7.9039	7.9019	7.8998	7.8978	7.8957
130	7.2961	7.2942	7.2923	7.2904	7.2885
140	6.7742	6.7725	6.7707	6.7690	6.7672
150	6.3234	6.3217	6.3201	6.3184	6.3168
160	5.9275	5.9259	5.9244	5.9228	5.9213

Table 3: Specific Absorption Rate (W/kg) at different depths in blood at different distances due to radiation from All India Radio Antenna of power 10 kW at 102.1 MHz

Distance (m)	10 μm	20 μm	30 μm	40 μm	50 μm
10	10.4763	10.4724	10.4685	10.4645	10.4606
20	2.6191	2.6181	2.6171	2.6161	2.6152
30	1.1640	1.1636	1.1631	1.1627	1.1623
40	0.6548	0.6546	0.6543	0.6541	0.6538
50	0.4191	0.4189	0.4187	0.4186	0.4184
60	0.2910	0.2909	0.2908	0.2907	0.2906
70	0.2138	0.2137	0.2136	0.2136	0.2135
80	0.1637	0.1636	0.1636	0.1635	0.1635
90	0.1293	0.1293	0.1292	0.1292	0.1291
100	0.1048	0.1047	0.1047	0.1046	0.1046
110	0.0866	0.0866	0.0865	0.0865	0.0865
120	0.0728	0.0727	0.0727	0.0727	0.0726

Table 4: Specific Absorption Rate (W/kg) at different depths in cerebro spinal fluid at different distances due to radiation from All India Radio Antenna of power 10 kW at 102.1 MHz

Distance (m)	10 μm	20 μm	30 μm	40 μm	50 μm
10	18.8526	18.8428	18.8330	18.8232	18.8134
20	4.7131	4.7107	4.7082	4.7058	4.7034
30	2.0947	2.0936	2.0925	2.0914	2.0903
40	1.1783	1.1777	1.1771	1.1765	1.1759
50	0.7541	0.7537	0.7533	0.7529	0.7525
60	0.5237	0.5234	0.5232	0.5229	0.5226
70	0.3848	0.3846	0.3844	0.3842	0.3840
80	0.2946	0.2944	0.2943	0.2941	0.2940
90	0.2327	0.2326	0.2325	0.2324	0.2323
100	0.1885	0.1884	0.1883	0.1882	0.1881
110	0.1558	0.1557	0.1557	0.1556	0.1555
120	0.1309	0.1309	0.1308	0.1307	0.1307
130	0.1116	0.1115	0.1114	0.1114	0.1113
140	0.0962	0.0961	0.0961	0.0960	0.0960
150	0.0838	0.0838	0.0837	0.0837	0.0836
160	0.0736	0.0736	0.0736	0.0735	0.0735

RESULTS AND DISCUSSION

Table 1 and 2 show the induced electric field and table 3 and 4 represent SAR at different distances from the broadcasting antenna in the blood and cerebro spinal fluid. The depth of penetration ranges from 10 μm to 50 μm . The broadcasting antenna of power 10 kW is used for calculation which is running at radio frequency i.e. 102.1 MHz to broadcast the signals. The values of SAR are above up to the distance 110 m than the value recommended by WHO and ICNIRP guidelines. However, the corresponding values for blood at a distance of 120 m are below the guidelines but, Table 4 represents that the corresponding values for cerebro spinal fluid up to a distance of 150 m are very near to the limiting point of the guidelines. These tables also represent the variation in induced electric field and SAR on increasing the depth inside the tissues. For blood the variation is not very large after 70 m and it become approximately constant after 120 m and for cerebro spinal fluid the variation is not very

large after 90 m and it become approximately constant after 160 m.

CONCLUSION

However, while technologies have been a vital part of our daily and we cannot avoid this, nor would we wish to, because technology makes our lives healthier, wealthier and safer but these radiations penetrate the tissues and may heat the tissue due to long term exposure, to its extent to bear which would be harmful for our health. Therefore, the international guidelines for installation of high power radio frequency transmitters must be made very clear and followed strictly. The installation of high power broadcasting antenna must be kept atleast 150 m away from the high populated area. The public should also be made aware of the use of radio frequency sources and the minimum distance which must be kept by them to live safer and healthy.

REFERENCES

- Dein, A.Z.E. and Amr, A., 2010. Specific Absorption Rate (SAR) Induced in Human Heads of Various Sizes When Using Mobile Phone, *Proceeding of the World Congress on Engineering (WCE), London (U.K.)*, **1**, June 30 – July 2.
- Gabriel, C., Gabriel, S. and Corthout, E., 1996. The Dielectric Properties of Biological Tissues: I. Literature Survey, *Phys. Med. Biol.* **41**, 2231-2249.
- Gabriel, S., Lau, R.W. and Gabriel, C., 1996. The Dielectric Properties of Biological Tissues: II. Measurements in the Frequency Range 10 Hz to 20 GHz, *Phys. Med. Biol.* **41**, 2251-2269.
- Gabriel, S., Lau R.W. and Gabriel C., 1996. The Dielectric Properties of Biological Tissues: III. Parametric Models for the Dielectric Spectrum of Tissues, *Phys. Med. Biol.* **41**, 2271-2293.
- Gandhi, F.M. and Mohammad, M.A.S., 2008. Thermal Effects of Radiofrequency Electromagnetic Fields on Human Body, *Journal of Mobile Communication*, **2(2)**, 39-45.
- Hardell, L., Carlberg, M., Ohlson, C.G., Westberg, H., Eriksson, M. and Hansson, M.K., 2007. Use of Cellular and Cordless Telephones and Risk of Testicular Cancer, *Int J Androl*, **30(2)**, 115-122.
- Hirata, A., Shirai, K. and Fujiwara, O., 2008. On Averaging Mass of SAR Correlating with Temperature Elevation Due to a Dipole Antenna, *Progress in Electromagnetics Research, PIER*, **84**, 221-237.
- Johansen, C., 2004. Electromagnetic Fields and Health Effects – Epidemiologic Studies of Cancer, Diseases of the Central Nervous System and Arrhythmia – Related Heart Disease, *Scand J Work Environ Health*, **30 (Suppl 1)**, 1-30.
- Joseph, W. and Martens, L., 2006. Reconstruction of the Polarization Ellipse of the EM Field of Telecommunication and Broadcast Antennas by a Fast and Low-Cost Measurement Method, *IEEE Trans Electromagn. Compat.*, **48 (2)**, 385-396.
- Kumar, V., Vats, R.P. and Pathak, P.P., 2008. Harmful effects of 41 and 202 MHz radiations on some body parts and tissues, *Indian Journal of Biochemistry & Biophysics*, **45**, 269-274.
- Nielsen, J.B., Elstein, A., Hansen, D.G., Kildemoes, H.W., Kristiansen I.S. and Stovring, H., 2010. Effect of Alternative Styles of Risk Information on EMF Risk Perception, *Bioelectromagnetics*, **31**, 504-512.
- Osepchuk, J.M. and Petersen, R.C., 2003. Historical Review of RF Exposure Standards and the International Committee on Electromagnetic Safety (ICES), *Bioelectromagnetics Suppl.*, **6**, S7-S16.
- Osepchuk, J.M., and Petersen, R.C., 2008. Safety and Environmental Issues, In: RF and Microwave Applications and Systems, Taylor & Francis Group, LLC, pp: 21.1-21.21.
- Ozen, S., Helnel, S. and Cerezci, O., 2008. Heat Analysis of Biological Tissue Exposed to Microwave by Using Thermal Wave Model of Heat Transfer (TWMBT), *Burns*, **34**, 45-49.
- Pathak, P.P., Kumar, V. and Vats, R.P., 2003. Harmful Electromagnetic Environment Near Transmission Tower, *Indian J. Radio Space Phys.*, **32**, 238-241.
- Prasad, K.D., 1999. Electromagnetic Waves, In: Electromagnetic Fields and Waves, I st ed, India: Satya Prakashan, pp: 425-520.
- Sirav, B. and Seyhan, N., 2009. Radio Frequency Radiation (RFR) from TV and Radio Transmitters at a Pilot Region in Turkey, *Radiat. Prot. Dosimetry*, **136 (2)**, 114-117.
- Stuchly, M.A., and Stuchly, S.S., 1996. Experimental Radio Wave and Microwave Dosimetry, In: Handbook of Biological Effects of Electromagnetic Fields, Eds, C. Polk and E. Postow, 2 nd ed, Boca Raton, CRC Press, pp: 295-336.

- Takebayashi, T., Akiba, S., Kikuchi, Y., Taki, M., Wake, K., Watanabe, S. and Yamaguchi, N., 2006. Mobile Phone Use and Acoustic Neuroma Risk in Japan, *J Occup Environ Med*, **63(12)**, 802-807.
- Takebayashi, T., Varsier, N., Kikuchi, Y., Wake K., Taki, M., Watanabe, S., Akiba S. and Yamaguchi, N., 2008. Mobile Phone Use, Exposure to Radiofrequency Electromagnetic Field and Brain Tumour: A Case-Control Study, *Br J Cancer*, **98**, 652-659.
- Wessapan, T., Srisawatdhisukul, S., Rattanadecho, P., 2011. The Effects of Dielectric Shield on Specific Absorption Rate and Heat Transfer in the Human Body Exposed to Leakage Microwave Energy, *International Communications in Heat and Mass Transfer* **38**, 255–262.
- Wideman, P.M., and Schutz, H., 2008. Informing the Public About Information and Participation Strategies in the Siting of Mobile Communication Base Stations: An Experimental Study, *Health Risk Soc.*, **10**, 517-534.

