

Vol. 10. No.4. 2021

©Copyright by CRDEEP Journals. All Rights Reserved.

Contents available at:

www.crdeepjournal.org

International Journal of Environmental Sciences (ISSN: 2277-1948) (CIF: 3.654)

A Peer Reviewed Quarterly Journal



Full Length Research Paper

Fish Species Diversity of the River Narmada in Nimar Region of Madhya Pradesh, India.

Khichi Yogesh

Head of the Department of Zoology Choithram College Indore, Madhya Pradesh, India.

ARTICLE INFORMATION

Corresponding Author:

Khichi Yogesh

Article history:

Received: 18-10-2021

Revised: 25-10-2021

Accepted: 29-10-2021

Published: 01-11-2021

Key words:

Fish Diversity, Fresh Water Fishes, Narmada River, Conservation

ABSTRACT

The fishes are one of the most important vertebrate, provided rich protein sources for human and several animals and important elements in the economy of many countries. Fish diversity of river essentially represents the fish faunal diversity. Rivers conserve a rich variety of fish species diversity monitoring of Nimar region carried out for one year i.e., August 2015 to July 2016. The aim of the study was to explore the fish fauna of Narmada River. In the course of investigation, Four sampling spots were selected i.e., Omkareshwar, Mandleshwar, Maheshwar and Barwani of Narmada River. The total 43 fish species were recorded under four orders, nine families and 16 genera. 21 species of commercial fisheries. The country is rich in diversity of such important group of animals Keeping in the view, the diversity of fish fauna of the Narmada River in Omkareshwar to Barwani, Madhya Pradesh, Central India has been studied Cypriniformes, three species of Ophiocephaliformes, two species. The Cyprinidae family is dominant and sub dominant family is of Perciformes and one species of Mastacembeliformes have been recorded Ophiocephalidae. Recently, the Tor tor fish species commonly called 'Mahasheer' has been declared state Fish of Madhya Pradesh. The over fishing and pollution are the major threat for fish diversity of the Narmada River.

Introduction

Study of biodiversity of fish fauna and their identification is one of the interesting field of biological research, which gives us an idea about the morphological variations and population diversity of fauna in polluted and non polluted site of any particular habitat (Napit K.M. 2013). Rich biodiversity of any ecosystem is absolutely essential in order to maintain their stability for proper function of their food chains (Siddiqui *et.al.* 2014). Fishes are the important element in the economy of many nations as they have been as table in the diet of many people (Shukla. P. *et.al.* 2013). Ichthyofaunal documentation is important to analyze status of fish species and also helps us for future planning to improve and conserve the biodiversity (Bose A.K. *et. al.* 2013). Present study is based on the fish diversity of Narmada River at Nimar region. The Narmada is a river in Central India and fifth largest river in subcontinent. It forms the traditional boundary between North India and South India. Narmada "the backbone of Madhya Pradesh" is the largest westward flowing river of India It is also reffered as 'lifeline of Madhya Pradesh'. It is considered holy by Hindus. It originates from Maikal Hill, Amarkantak in Shahdol district of Madhya Pradesh. It is situated at longitude 72 32' and 81 45'E and latitude 21 20' and 23 45'N. Total length of River Narmada is 1312 km which after traveling through three states namely Madhya Pradesh, Maharashtra and Gujarat for a distance of 1,077 km, 74 km and 161 km, respectively joins the Gulf of Cambay, near the District of Bharuch, Gujarat. (Pathak T. *et.al.*2014). The fish are a major food source for people, the study of ichthyology also has economic importance. This may have been because fish were both a relatively easy-to-obtain source of food as well as an animal group that was readily accessible, since fishing is one of humankind's oldest occupations.

Material and method

Description of Study Area

The Narmada river is considered as the life line of Madhya Pradesh. The catchment area of the river exists in the States of Madhya Pradesh (86.18%), Gujarat (11.6%), Maharashtra (1.5%), and Chattisgarh (0.72%). During its course, the river drops from an elevation of 1051 m to sea level, and flows through narrow gorges in the head reaches. The basin is bounded on the north by the Vindhya ranges, on the east by the Maikal range, on the south by the Satpura ranges and on the west by the Arabian Sea. Deep black soil covers the major portion of the basin. The river has 41 tributaries, of which 22 are on the left bank and 19 are on the right bank. The Barna,

Tawa, Kolar, and Sukta dams have been constructed on the tributaries. The Bargi is constructed on the mainstream, while the Indira sagar, Omkareshwar, Maheshwar and Sardar Sarovar dams are under construction.




Fig. 1: Map of Narmada river showing sampling stations

Methodology

The sampling was carried out seasonally covering pre-monsoon, monsoon, post-monsoon and winter season. Experimental fishing was carried out with the help of local fishers. After obtaining the fish from the site photographs were taken. The specimens were taken to the laboratory for identification. Morphometric measurements were taken and meristic characters were observed and the fin formula was completed. The fish samples were preserved in 5-10% formalin according to the size of the fishes. Smaller fishes were directly placed in the formalin solution, while larger fishes were given an incision on the abdomen before they were fixed. Plastic jar were used for the collection and preservation. Fishes were labeled based on the serial number, common name, scientific name, locality and date of collection. Fishes were identified with the help of taxonomic key, Days fauna (1994) and Talwar and Jhingran 1991. Fish Base website was also referred for various aspects of fish fauna (www.fishbase.org). Specific identifying characters on the body was observed and noted. Omkareshwar, to Barwani Flowing of Narmada river. Narmada Nimar area is **considered the mother and giver of peace**. The river basin is home to India's best teak and hardwood forests. Large irrigation projects have been completed on Narmada River, which supply water to hundreds of farmers all over central India.

Results and discussion

A total 43 species of fishes recorded from selected sites of Narmada River at nimar region belonging to 7 orders and 14 families. Among species, family Cyprinidae was the most dominant with 24 species and the percentage composition is 59% of fishes followed by Bagridae 9% with 4 species, Siluridae and Ophiocephalidae 7% with 3 species, Claridae, Notopteridae and Mastacembellidae 4% with 2 species and Schielbeidae, Clupeidae, Belonidae, Centropomidae, Nandidae, Gobioidae and Anabantidae represented with single species of fish in each family with 2% of each.

The species diversity peak is in post monsoon coinciding with favorable conditions such as sufficient water and ample food resources. The diversity was low in pre monsoon probably due to the shrinkage of water. Information collected from fisherman communities displayed high decline of fish diversity. Deforestation, water scarcity, pollution, introduction of exotic species and mining and excessive fishing are the biggest threats to fish population. Various workers have done work on Narmada river. Vishwakarma *et.al.* (2014), recorded 33 fish species belonging to 5 orders, 9 families and 21 genera.

Kumar *et.al.* (2014) studied the fish species diversity of river Narmada In Khedighat, Warwaha, Madhya Pradesh, and recorded 21 species of fish belonging to 4 orders and 6 families. Family cypriniformes were dominated with 15 species of fish.


Pathak *et.al.* (2014) recorded 58 species of fish from western region of Narmada river at nimar region. Vyas *et.al.* (2013) recorded 27 species of fish from Jamner river, a tributary of Narmada river.

Siddiqui *et.al.*(2014) work done on Biodiversity of Ichthyofauna of Narmada river of Mandleshwar region, Madhya Pradesh, India and recorded 48 species of fish belonging to 7 orders and 17 families. Bose *at.al.* (2013) recorded 57 species, belonging to 35 genera , 13 families, and 6 orders from middle stretch of river Tawa.

Bakawale *et. al.* (2013) worked on the fish Species diversity of the River Narmada in western zone, and recorded total 51 species of fish belonging to 7 orders and 15 families. In the present study 49 fish species, belonging to 7 orders and 14 families were recorded.

Present investigation revealed that, Narmada river is a healthy water body providing a habitat for freshwater fishes of diverse type. However, there is constant threat to fish population due to eutrophication and illegal fishing activities. The illegal fishing activities should be banned to prevent depletion of fresh water fish resources and further studies should be conducted to generate more details

regarding seasonal production and ecology of fishes. In situ conservation is one of the several prominent and suggestive measures for the conservation of fish biodiversity.



Cyprinidae (51%) > Bagridae (9%) > Siluridae(7%) > Ophiocephalidae(7%) > Clariidae (4%) >Notopteridae (4%) > Mastacembelidae(4%) > Scheilbedae (2%) ,Clupeidae (2%), Belonidae(2%),Centropomidae (2%), Nandidae (2%), Gobiidae (2%), Anabantidae (2%).

Fig. 2 % Composition of fish families in Narmada river during Aug. 2015 to July. 2016.

Table 1 : List of fish species from Narmada river

Order	Family	species
Cypriniformes	Cyprinidae	Labeo rohita
		Catla catla
		Cirrihinus cirrhosa
		Cirrihinus reba
		Labeo calbasu
		Labeo fimbriatus
		Labeo bata
		Labeo gonius
		Nemacheilus botia
		Puntius chola
		Puntius sarana
		Puntius dorsalis
		Tor tor
		Ctenopharygdon idella
		Hypothalmichthys molitrix
		Cyprinus carpio
		Puntius ticto
		Rasbora daniconius
		Siluriformes
Heteropneustes fossilis		
Hilsa hilsa		
Ompok bimaculatus		
Wallago attu		
Siluridae	Mystus bleekeri	
	Mystus seenghala	
Bagridae	Mystus cavasius	
	Mystus aor	
	Mystus tangara	
	Mystus vittatis	
	Rita rita	
Cluiformes	Clupeidae	Notopterusnotopterus
		Notopterus chitala
Beloniformes	Belonidae	Xenentodon cancila
Perciformes	Centropomidae	Chanda nama
		Nandus nandus
	Anabantidae	Anabas testudinus
Ophiocephaliformes	Ophiocephalidae	Channa marulius
		Channa punctatus
		Channa striatus
		Channa gachua
Mastacembeliformes	Mastacembelidae	Mastacembelus armatus
		Mastacembelus pancalus

Conclusion: Narmada river in nimar region is a important fresh water of Omkareshwar to Barwani region it is useful for Commercially important for fishes species diversity. The water of Narmada river is ideally, the quality wise suitable for fish culture. There is 43 species identified.

References

- Bakawale and Kanhere R. R. (2013). Study on the Fish Species Diversity of the River Narmada in Western Zone. *Research Journal of Animal, Veterinary and Fishery Sciences*,1(6):18-20.
- Bose A.K., B.C. Jha, V.R. Suresh, A.K. Das, A. Parashar and Ridhi.(2013).Fishes of the middle stretch of river Tawa, Madhya Pradesh, India. *Journal of Chemical, Biological and Physical sciences*.vol 3(1): 706-716.
- Day Francis, F.L.S. and F.Z. S. *The fishes of India* (1994), Jagmander book agency, New Delhi,vol 1-2
- Napit Mukesh kumar,(2013). Study of fish fauna of Bundelkhand region with special reference to Damoh district. *International journal of advance research*. Vol 1(4):24-30.
- Pathak , K. Borana & T. Zafar,(2014),Ichthyofauna of western region of Narmada river, Madhya Pradesh. *International Journal of Research in Applied ,Natural and SocialSciences*, Vol. 2, Issue 4, 25-28.
- Paunikar , S.S. Jadhav and S.S Talmale, (2012). Studies on Ichthyofaunal Diversity of Gour River, Jabalpur, Madhya Pradesh, Central India. *World Journal of Fish and Marine Sciences* 4 (4): 356-359.
- Shukla P and Ajay Singh (2013), Distribution and Diversity of Freshwater Fishes in Aami River, Gorakhpur, India. *Advances in Biological Research* 7 (2): 26-31.
- Siddiqui Anis, Meenakshi Chouhan, and Shailendra Sharma (2014), Biodiversity of Ichthyofauna of Narmada River of Mandleshwar Region, Madhya Pradesh, India, (2014). *Science Secure Journal of Environmental Biology*. Vol 1(1), 21-25.
- Talwar P.K. and A.G. Jhingran (1991). *Inland fishes of india and adjacent countries*. Vol 1and 2. Oxford and IBH publishing co. pvt. Ltd. New Delhi India.
- Vishwakarma Kripal Singh, Altaf Ali Mir, Abhilasha Bhawsar and Vipin Vyas, (2014) Assessment of Fish assemblage and distribution in Barna Stream Network in Narmada basin(Central India), *International Journal of Advanced Research*, Volume 2(1), 888-897.
www.fishbase.org.
- Yodha Ravindra Kumar and Chaurasia Rajendra Kumar, (2014).Studies on the fish species diversity of river Narmada in Khedighat, Barwaha, MP, India. *International Journal of Developmental Research and Engineering*. 1(1).