

**Full Length Research Paper**

Study on assessment of water quality of Punpun River at selected places of Aurangabad District, Bihar, India

¹*Dr Arvind Kumar Rai and ²Dr Avinash Kumar

¹Garhwa Polytechnic College, Garhwa, Jharkhand, India.

²GGSES, Bokaro, Jharkhand, India.

ARTICLE INFORMATION**ABSTRACT****Corresponding Author:**

Arvind Kumar Rai

Article history:

Received: 03-06-2022

Revised: 10-06-2022

Accepted: 11-07-2022

Published: 16-07-2022

The present research paper work focuses on the variations in different Punpun river water quality parameters like of the selected locations at Siris marked as S_1 , Santhua marked as S_2 & Jamhor S_3 village in Aurangabad District, Bihar. The water samples collected from the selected sites were analyzed at specific time intervals. The parameters studied were namely Temperature, pH, Dissolved Oxygen (DO), Total Suspended Solids (TSS), Total dissolved Solids (TDS), Alkalinity, Hardness and Chloride. This study also helps us to understand the changes in water quality due to anthropogenic influence. However, no attempt has been made to analyze the water quality of these areas.

Key words:

Punpun, Water quality, pH, Hardness, Chloride.

Introduction

It is well known fact that clean water is absolutely essential for healthy living. Freshwater resources all over the world are threatened not only by over exploitation and poor management but also by ecological degradation. The main source of freshwater pollution can be attributed to discharge of untreated waste, dumping of industrial effluent, and run-off from agricultural fields. Industrial growth, urbanization and the increasing use of synthetic organic substances have serious and adverse impacts on freshwater bodies. Freshwater resources are under severe and increasing environmental stress. (US EPA, 2016.) The problem of fresh water pollution in India came to the forefront towards the beginning of 1960 with the domestic waste water, sewage, effluent and industrial waste water discharges being the most critical sources of pollution in various towns and cities. In recent years, because of continuous growth in population, rapid industrialization and various technologies involving waste disposal (Jain, P, 2009). Human factor is the main cause of river pollution.

In Aurangabad, Punpun is the primary source of drinking water. In the last few decades, there has been a serious concern over the deterioration in its water quality. The river has been receiving large amounts of be indiscriminate disposal of domestic effluents, immersion of idols, dumping of plastics into drain, bathing of cattle and agricultural runoff, partially treated and untreated wastewater during its course, especially between Tandwa to Daudnagar areas. Pollutants flowing into the river are contributed from the waste of the villages situated along its bank. Table No 1 gives the drinking water quality standards as per IS: 10500-1991.

Table 1. Drinking Water Quality Standards (IS: 10500-1991)

S. No	Parameters	Range	Effects
Physical			
1	Turbidity, Units on J.T.U scale	2.5	Undesirable effects on human beings
2	Color	5.0	Undesirable effects on human beings

3	Taste and Odor, Units on Platinum cobalt scale	Unobjectionable	Undesirable effects on human beings
Chemical			
4	pH	6.5-8.5	Beyond this range the water will affect water supply system and mucous membrane
5	Total hardness	300	Adverse effect on domestic use
6	Iron	0.3	Adverse effect on human health
7	Chloride	250	Cause Taste, corrosion and palatability are affected
8	Residual free chlorine	0.2	--
9	Dissolved solids	500	Cause gastrointestinal irritation
10	Nitrates	45	Beyond this Blue Baby Syndrome may be caused in infants.
11	Fluoride	1.0	Cause fluorosis
12	Alkalinity	200	Beyond this water becomes unpleasant
13	PAH, Hexavalent chromium	0.01	May be carcinogenic
14	Pesticides	Absent	Very toxic
15	Heavy metals	0.001	Becomes more toxic

Materials and Methods

Study area

It has an average elevation of 138 metres (452 feet) and 35 km from Aurangabad. Figure No 1 gives the map of study area. The three sampling stations Siris (S_1), Santhua (S_2) and Jamhor (S_3) were marked out for the collection of water samples. The Punpun River is a sacred river for Hindus. The 210 km long river is mostly rainfed and carries little water in the dry season. The Punpun River is a tributary of the Ganges. It originates in Gajandham, Palamu District of Jharkhand. It flows through Chatra, Aurangabad, Gaya and Patna Districts of Jharkhand and Bihar. The river mostly flows in a north east direction and the join the Ganges at Fatuha, Bakhtiyarpur, Patna. The main tributaries of the Punpun are the Adari, Batane, Batre, Madar and Morhar. The catchment area of the Punpun is 8,530 square kilometers. Basic information of Aurangabad District and study area is shown in Table No 2.




Fig 1. Map showing study area

Collection of samples

Water samples were collected Siris, Santhua and Jamhor village (8.30 AM) from the locations along the route of the Punpun River from different locations at Aurangabad areas. The water samples were collected in the cleaned, air tight water glass bottle and taken to the laboratory for analysis as per guidelines of APHA, 1995. The water samples were transported under ice cooled conditions with appropriate sealing.

Results and Discussion

Analysis of the selected parameters of the Punpun River is shown in Table No 3.

Table 3. Physico Chemical Water Quality Parameters of Punpun

S. No	Parameters /units	Methods	S1	S2	S3
			Average	Average	Average
1	Temperature, 0°C		27.54	29.40	22.21
2	pH	Potentiometer	7.31	7.52	7.0
3	DO, mg/l	Winkler's method	5.24	6.40	5.29

4	TSS, mg/l	Gravimetric method	118.40	128.73	123.26
5	TDS, mg/l	Gravimetric method	238.23	243.54	229.14
6	Alkalinity, mg/l	Titration	40.12	28.40	33.56
7	Hardness, mg/l	EDTA method	112.27	65.27	83.31
8	Chloride, mg/l	Titration	30.25	34.56	33.61

Temperature

The temperature of water has very important ecological effects. Arise in temperature of the water leads to the speeding up of the chemical reactions in water, reduces the solubility of gases and amplifies the tastes and odours. Water in the temperature range of 7⁰C to 11⁰C has a pleasant taste and is refreshing. At higher temperature with less dissolved gases the water becomes tasteless and even does not quench the thirst. In present study, the values of temperature were ranged between 22.21⁰C to 29.40⁰C.

pH

pH is the one of the most important single factor which influences aquatic production. pH has no direct adverse effects on health, however a lower value below 4.5 will produce odd taste; and higher value above 7.5, an alkaline taste. Most rivers have a neutral pH of 6.5-8.5. Low pH levels (below optimal) can result in fish kills by stressing their systems causing physical damage, which in turn can make them more vulnerable to disease, similarly high pH particularly in combination with high water temperature, can increase the amount of unionized ammonia which is highly toxic to fish (Kumar et al.,2010). In present study, the values of pH were ranged from 7.0 to 7.52. All the pH values represent alkaline nature of river water samples were within the permissible limit as per IS: 10500.

Dissolved Oxygen

Oxygen saturation in the environment refers to the amount of oxygen dissolved in the water bodies. Water bodies have different types of micro-organisms such as virus, bacteria, and fungi. When too much biodegradable matter end up in water bodies, it encourages more microorganism growth, and they use up more oxygen in the water. If oxygen is depleted, aerobic organisms die, and anaerobic organism grow more to produce harmful toxins such as ammonia and sulfides (Mishra et al., 2009). In the present study, the values of DO were ranged from 5.24 mg/l to 6.40 mg/l.

Total Suspended Solids (TSS)

Total suspended solids are particles that are larger than 2 microns found in the water column. Anything smaller than 2 microns is considered a dissolved solid. Most suspended solids are made up of inorganic materials, though bacteria and algae can also contribute to the total solids concentration. In present investigation, TSS was ranged from 118.40 mg/l to 128.73 mg/l.

Total Dissolved Solids (TDS)

In natural waters dissolved solids are composed mainly of carbonates, bicarbonates, chlorides, sulphates, phosphates, and nitrates of calcium, magnesium, sodium, potassium, iron and manganese etc. In present investigation, TDS was ranged from 229.14 to 243.54 mg/l. All the TDS values of river water samples were within the permissible limit as per IS: 10500.

Alkalinity

Alkalinity is also important to aquatic organisms because it protects them against rapid changes in pH. During the study period, Alkalinity values ranged between 28.40 mg/l to 40.12 mg/l. Alkalinity itself is not harmful to human beings but desirable limit of 100 mg/lit is always required for domestic supply of water.

Hardness

The total hardness of water is defined as the sum of calcium and magnesium concentrations, both expressed as calcium carbonate, in mg/L. Based on present investigation, hardness varied from 65.27 mg/l to 112.27 mg/l. The permissible limit of Hardness for drinking water is 300 mg/l (IS 10500). Hence, all the hardness values of river water samples in study area were within the permissible limit.

Chloride

Chloride may get into surface water from several sources such as industries, municipalities, agricultural runoff and other sources. During the study period, Chloride varied from 30.25 to 34.56 mg/l. Hence, all the Chloride values of Punpun river water samples were within the permissible limit. The EPA Secondary Drinking Water Regulations recommend a maximum concentration of 250 mg/litre for chloride ions.

Conclusion

The above mentioned results show severe deterioration of water quality of the Punpun River at selected areas of Aurangabad. Domestic waste water, agricultural run-off, urban effluents, fertilizer run-off contain phosphorus and nitrogen, manure from

livestock operations, which increase the level of nutrients in water bodies and can cause deterioration of water quality in the rivers. The nitrates come mainly from the fertilizer that is added to the fields. Excessive use of fertilizers cause nitrate contamination of groundwater, with the result that nitrate levels in drinking water is far above the safety levels recommended. Organic matter and nutrients causes an increase in aerobic algae and depletes oxygen from the water column. This causes the suffocation of fish and other aquatic organisms in and around Aurangabad district.

References

- A. P. H. A. 1995. Standard Methods for the Examination of Water and Waste Water 19th Edition, New York, USA.
IS: 10500-1991: Indian Standards for Drinking Water Quality as per BIS specifications.
Jain, P. 2009. Sick Yamuna, Sick Delhi-Searching Correlation. Peace Institute Charitable Trust.
Kumar, A., Bisht, B.S., Singh, A.K., Talwar, A.2010. Physical, Chemical and Bacteriological Study of Water from Rivers of Uttarakhand, *Journal of Human Ecology*, 32(3): pp169-173.
Mishra, A., Mukherjee, A. and Tripathi, B.D. 2009. Seasonal and Temporal Variation in Physico Chemical and Bacteriological Characteristics of River Ganga in Varanasi, *International Journal of Environmental Research*, 3(3): 395- 402.
US EPA 2016. Water Quality Assessment and TMDL information. Washington, DC, United States Environmental Protection Agency (US EPA)