



Content is available at: CRDEEP Journals
Journal homepage: <http://www.crdeepjournal.org/category/journals/ijs/>

International Journal of Life Sciences
(ISSN: 2277-193x) (Scientific Journal Impact Factor: 6.106)

UGC Approved-A Peer Reviewed Quarterly Journal



Full Length Research Paper

Studies on Limnological Characteristics of Sowlanga New Tank, Nyamathi Davangere Karnataka India

Prabhavathi S. Horadi¹

Assistant Professor & H.O.D of Botany, A.V.K College for Women Davangere, Karnataka-577004, India.

ARTICLE DETAILS

Corresponding Author:
Prabhavathi S. Horadi

Key words:

Ramankere tank,
Physion-chemical
parameters, Planktonic
composition

ABSTRACT

An investigation was carried out in characteristics but in Sowlanga tank near Nyamathi of Davangere on physico-chemical and Planktonic composition during January to parameters were compared with the December 2024. The results of physico-chemical standard values prescribed by the Bureau of India Standards (BIS) and World Health Organization (WHO). The study revealed that, tank water is polluted as it possesses high BOD, CO₂, phosphate and nitrogen. A total of 36 species belonging to 27 genera of phytoplanktons were recorded, of which Bacillariophyceae and chlorophyceae were dominant. Eight zooplanktons and six macrophytes were also recorded. In the light of standard of water quality recommended by WHO, the tank water should not be used by human beings especially for drinking and cooking.

1. Introduction

Water is renewable natural resources essential for all life sustaining processes on earth Majority of water available on earth his saline in nature: only a small quantity exists as fresh water. water Fresh water has become a scarce commodity due to over exploitation and its necessities have led to the deterioration of surface and subsurface water. The causative factors for the pollution of water are industries, agriculture and domestic activities (Prabhakaran et al., 2011). Due to over expanding population and industrial settlements, the demand for fresh water is increasing day by day. In today's scenario, unplanned urbanization, rapid industrialization and indiscriminate use of artificial chemicals cause heavy and varied pollution in aquatic environments leading to deterioration of water quality and depletion of aquatic fauna. Physico-chemical parameters play a vital role in determining the distributional pattern and quantitative abundance of organisms inhabiting a particular.

Physico-chemical parameters play a vital role in determining the distributional pattern and quantitative abundance of organisms inhabiting a particular aquatic ecosystem (Santhoshkumar Singh et al., 2009). Several investigators have studied the physico-chemical dynamics of varied lentic water bodies with the intent to assess the water quality (Sayeswara et al., 2010; Purushothama et al., 2011; Sayeswara et al., 2011; Sayeswara et al., 2011a; Mahesh and Sayeswara, 2011; Mahesh et al., 2012) Planktons form the base of food chain in aquatic ecosystems, thus playing a vital role in fisheries. The spatial and temporal variation of planktons is regulated by major environmental factors. Several studies have been done on the phytoplankton diversity at ponds and tanks (Nafessa et al., 2010; Nafeesa et al., 2011; Purushothama et al., 2011a; Sayeswara et al., 2012; Sayeswara et al., 2012a; Sayeswara et al., 2012b; Vasantha et al., 2013). The plankton study is a very useful tool for the assessment of water quality, trophic status and pollution level. Ramankere tank is an annual water body receiving water from the adjacent paddy fields, Tunga irrigation canal. The total area of Ramankere tank is about 85 acres of which water spreads over an area of 270 acres with an average depth of 20 feet. It is located at Sowlanga, 20 km away from Shivamogga town. The water is used for domestic purposes like washing of clothes, vehicles and for domestic animals, etc. The water has undergone moderate changes in its physico-chemical properties due to ecological degradation,

¹Corresponding Author can be contacted at: Assistant Professor & H.O.D of Botany, A.V.K College for Women Davangere, Karnataka-577004, India.

Received: 04-Jan-2025; Sent for Review on: 18-Jan-2025; Draft sent to Author for corrections: 20-Jan-2025; Accepted on: 30-Jan-2025; Online Available from 12-Feb-2025

DOI: [10.13140/RG.2.2.10675.54567](https://doi.org/10.13140/RG.2.2.10675.54567)

IJLS-8799/© 2025 CRDEEP Journals. All Rights Reserved.

overflowing of water from adjacent paddy fields and other excessive human activities. The basis of selection of Sowlanga new tank was that its water is used by a large population which receives adequate waste water and periodic flooding from plains. In the present investigation, an attempt has been made to assess the plankton diversity and suitability of water for human consumption and domestic purposes.

2. Materials and methods

2.1 Study area :

Water was sampled on monthly basis, between 7 to 9 am from January to December 2024. This water samples were collected in good quality polythene bottles. Water temperature was recorded at the sampling site itself. Dissolved oxygen was fixed on the spot itself in BOD bottles. Various parameters like turbidity, total hardness, sulphate, free CO alkalinity, BOD, TDS, phosphate, nitrate and chloride were estimated as per the standard methods (APHA, 1998). Plankton samples were collected by using plankton net (NO.1) by filtering 100 liters of water, preserved in 4% formaldehyde. Identifications of phytoplanktons and zooplanktons were made with the help of Deshikachary (1959). Gandhi (1961), Welch (1952) and Prescott (1982).

3. Results and discussion

The results of seasonal variation of physico-chemical parameters of Sowlanga new tank are given in Table 1.

- *Temperature:* The water temperature is largely influenced by local climatic conditions. The seasonal water temperature ranged from 20.2 to 26.9°C. The minimum value recorded in December and maximum in April. Turbidity is a measure of cloudiness of water.
- *Turbidity:* Turbidity in natural water arises due to the presence of suspended matter such as clay, silts, finely divided organic and inorganic matter, phytoplanktons and other microscopic organisms. The values of turbidity ranged from 10.3 to 57.5 NTU. The highest and the lowest values were recorded in July and March, respectively.
- *pH:* pH values are slightly acidic to slightly alkaline and found within permissible limit of 6.3 to 7.4 as per the Bureau of Indian Standards (BIS, 1993). The minimum value observed during January (6.3) and maximum during July (7.4). The pH is important since aquatic organisms are well adapted to specific pH range and do not withstand abrupt changes in it.
- *Dissolved oxygen:* Dissolved oxygen is an important gaseous factor that determines the quality of water and intern regulates the distribution of aquatic organisms. In the present study the DO level fluctuated between 2.1 to 5.3 mg/L. The highest and the lowest values were recorded 1 in October and February, respectively. The variations of DO depend on the primary production and respiration of aquatic organisms. The permissible standard of DO is above 5mg/L (Perk and Park, 1980).

Table1. Physico-chemical characteristics of Sowlanga new tank water.

Parameters	Months-2024											
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Temperature	22.6	22.9	23.8	26.9	25.9	23.3	21.3	22.2	22.9	22.4	21.8	20.2
Turbidity	12.7	11.9	10.3	13.4	20.4	48.1	57.5	32.4	34.9	26.6	13.8	12.7
pH	6.3	6.5	6.4	7.3	7.1	6.8	7.4	7.2	7.1	7.2	7.3	7.2
DO	3.2	2.1	2.5	2.8	3.5	3.8	3.7	3.9	4.3	5.3	3.5	2.9
BOD	6.1	6.3	6.2	7.8	7.1	7.8	7.5	8.2	8.4	7.1	7.1	7.8
CO2	14.9	15.1	22.9	21.5	27.5	15.2	16.1	28.9	19.3	18.7	21.9	26.4
Alkalinity	74.3	78.4	63.8	70.6	72.1	62.7	60.4	61.1	60.3	51.2	54.7	61.7
TDS	33.6	34.5	40.9	57.3	40.7	40.1	41.1	41.1	33.8	31.3	41.4	40.4
TH	87	81	119	103	107	74	70	79	99	83	68	97
Chloride	72.1	70.1	80.2	81.4	82.9	72.1	64.4	71.1	62.2	78.1	79.3	72.3
Phosphate	1.1	1.2	1.5	1.6	1.6	1.2	1.2	1.3	1.2	1.6	2.3	1.7
Nitrate	5.3	4.7	6.2	6.1	7.8	6.9	6.1	6.5	5.2	5.1	4.2	3.7
Sulphate	11.3	10.7	9.5	7.1	10.1	8.8	8.4	7.4	8.7	12.1	7.3	7.6

- *Biological Oxygen Demand:* BOD is the measure of degradable organic matter present in water. BOD and other microbial activities generally increase by the introduction of sewage (Hynes, 1971). In the present study BOD values ranged between 6.1 to 8.4mg/L. The minimum value was noticed in the month of January while maximum in permissible limit of 6.5mg/L (WHO, 1991). They were found above the permissible limit of 6.5mg/L,(WHO-1991)
- *Carbon dioxide:* Free carbon dioxide values fluctuated between 14.9 to 28.9 mg/L. The highest and the lowest values of CO, was due to the absorption by plants for were recorded in August and January, respectively. The variation photosynthesis and activity of other living organisms.
- *Alkalinity:* Alkalinity in the water samples is primarily a function of carbonate, bicarbonate and hydroxide content. In the present study total alkalinity ranged from 51.2mg/L (October) to 78.4mg/L (February). It is within permissible limit of 600mg/L (BIS, 1993). Surface alkalinity may result from the discharge of domestic wastes.

- **Total Dissolved Solids:** TDS values ranged from 31.3 to 57.3mg/L, the minimum was recorded in October and maximum in February. The minimum value may be due to the stagnant condition of the water body. The values are within permissible limits of 1500 mg/L (BIS, 1993).
- **Total Hardness:** Total hardness of water is not a pollution parameter but indicates water quality mainly in terms of Ca² and Mg² contents. Total hardness values observed are 68 to 107mg/L. The minimum value was recorded in November and maximum in May.
- **Chloride:** Chloride is an important anion found in variable amounts in water bodies. Chlorides increase the degree of eutrophication (Goel et al., 1980). In the present study, chloride values fluctuated between 62.2mg/L (September) to 82.9 mg/L (May). High chloride content indicates the deterioration of water quality usually linked with sewage load (Mini et al., 2003). The most important sources of chlorides in the fresh water are the discharge of domestic sewage and farm drainage. The concentration of chlorides is thus the indicator of pollution.
- **Phosphate:** Phosphorus occurs in natural water as various types of phosphates. The most important source of phosphates are the discharge of domestic sewage, detergents and agricultural runoff. Values of phosphates ranged from 1.1 to 1.7 mg/L with the minimum value in January and maximum in December.
- **Nitrates:** Most of the unpolluted sources of water are deficient of nitrates because it exists only in few natural sources (Trivedi and Goel, 1984). In the present study, nitrate values ranged from 3.7 to 7.8mg/L. The minimum value of nitrate was noticed in the month of December while maximum in May.
- **Sulphate:** Sulphate is one of the major anions occurring in natural waters. It may enter natural waters through weathering of sulphate bearing deposits. The values fluctuated between 7.1 to 11.3mg/L. The minimum value was recorded in July and maximum is January. The relationship between various physico-chemical parameters of water samples were analyzed statistically conducting the Pearson correlation analysis (Table 3). Correlation analysis is an important part of bivariate analysis which is concerned with the relation between two variables.

3.1 Planktonic composition:

A total of 36 phytoplanktons, 08 zooplanktons and 06 macrophytes were identified, which are given in Table 2 & 3. Among phytoplanktons, members of Bacillariophyceae and Chlorophyceae appear to be dominant as compare to other classes. The acidic pH favors the abundance of chlorophycean members. The presence of Euglenophycean members indicate that the water is organically polluted as the Euglenoids are the bioindicators of pollution. Karatella and Daphni were noted among zooplanktons. Ipomea aquatica and Eichhornea crassipes were dominant among the macrophytes.

Table 2. List of Phytoplankton in Sowlanga newtank water.

Sl.no	Chlorophyceae		Euglenophyceae
1.	Coelastrum reticulatum	19	Eugleno gracile
2.	Crucigeniarectangularis	20	Eugleno elongata
3.	Eudorina legans	21	Phacuscurvicauda
4.	Oocystis gigas	22	Phacustruqueter
5.	Pediastrum simplex	23	Strombomonasgibberosa
6.	Scenedesmusdimorphon		Bacillariophyceae
7.	Selanastrum westii	24	Diatoma vulgare
8.	Tetraedonlongispinum	25	
9.	Tetraedon minimum	26	Gomphonemaabbreviatum
10.	Tetraedon muticum	27	Gomphonemalanceolatum
		28	Gyrosigmatenellum
	Cyanophyceae		
1.	Anacystis sp.	29	Melosiragranulata
2.	Gloecapsa sp.	30	Nitzchiaacicularis
3.	Merismopedia glauca	31	Pinnularia major
4.	Merismopedia tenuissima	32	Pinnularianobilis
5.	Nostoc microscopium	33	Suriellacapronii
6.	Oscillator formosa	34	Surielarobusta
7.	Phormidium sp.	35	Synedraacus
8.	Rivulera sp.	36	Synedra tabulate

The water samples from Ramankere tank was collected and analyzed for various physico-chemical parameters to study the extent of pollution. Planktonic composition was also studied. DO was very low and BOD, CO, phosphate and nitrogen values were significantly higher than the permissible level for domestic consumption. The presence of bioindicators of pollution (phytoplanktons and zooplanktons) indicates the occurrence of organic pollution. In the light of standard of water quality recommended by WHO, the tank water should not be used by human beings especially for drinking and cooking. Precautionary measures should be taken before the water is consumed. In order to maintain the health of the tank with respect to water quality it is essential that authorities should take immediate step on the following points.

- People should not be allowed to discharge domestic wastes directly in to the tank.
- Avoid washing of clothes.
- Awareness should be created regarding the impact of water pollution on the human health.
- People should be advised to avoid dumping of agricultural waste.

Table 3. List of Zooplanktons and macrophytes in Sowlanga new tank.

Cyclopes sp	05	Ipomea aquatica
Daphnia sp	06	Jussia repens
Keratella sp..	07	Nelumbo nucisera
Paramoecium caudata	08	Nympea nouchali
Macrophytes		
Azollapinnata	04	Salvinia natans
Cyperus sp	05	Trapa bipinosa
Ipomeaaquatica	06	Eichhorea crassipes

Acknowledgement: The author express their gratitude to Principal, AVK Collge for Women , Davangere for facilities and encouragement.

4. References

- APHA (1998). Standard Methods for the Examination of Water and Waste Water, 20th ed. Public Health Association, Washington, D.C.
- BIS: (1993). Methods of sampling and Test (Physical and Chemical) for water and waste water, Ist Revision, 1-2.
- Deshikachary, T. V. (1959). Cyanophyta, ICMR, New Delhi.
- Environmental Publications, Karad (India). Welch, P. C. (1952). Limnology. McGraw-Hill Publications, New York. WHO, (1991). International Standards for drinking water, Geneva.
- Gandhi, H.P. (1961). Notes on the diatomaceous of Ahmedabed and its environs. Hydrobiology, 17: 218-236.
- Goel, P. K, Gopal, B. and Trivedi, R.K. (1980). Impact of sewage on freshwater ecosystem, Int. J. Ecol and Environ. Sci., 6: 97-116.
- Hynes, H. B. N. (1971). The Biology of Polluted Water. Uni. Toronto Press, Canada.
- Mahesh Anand Goudar and Sayeswara, H.A. (2011). Hydrochemistry of Bhudhigere tank near Shivamogga, Karnataka, India. Current Biotica, 5(1): 85-90.
- Mahesh Anand Goudar, Sayeswara, H.A. and Goudarashivannanavar (2012). Assesment of tank water quality in relation of some physico-chemical parametrs-A case study in Abbalgere tank of Shivamogga, Karnataka, India. Asian Journal of Microbiology, Biotechnology and Environmental Science. 14(3): 385-389.
- Mini, I., Radhika, C. J.and Tunga Devi, T. (2003). Hydrobiological Thiruvananthapuram, Kerala, Poll. Res., 22(4): 617-626. studies on a Lotic Ecosystem, Vamanapuram River,
- Nafeesa Begum, Sayeswara, H.A. and Naik, K.L. (2011). Seasonal variations of Phytoplankton diversity in Bethur pond near Davangere, Karnataka, India. Environemnt & Ecology. 29(2A) : 1355-1357.
- Nafessa Begum, Narayana, J. and Sayeswara, H.A. (2010). A Seasonal study of Phytoplankton diversity and Pollution indicators of Bathi pond near Davangere city, Karnataka, India. Environment Conservation Journal (11(3): 75-80.
- Perk, J. E.and Park, K. E. (1980). A Text bok of preventives and social medicine, 8^a edition, Messer BanrsidasBhanot, Jabalpur.
- Prabhakaran, N., Mahendran, N., Radha, S., Gurugnanam, B. and Mahendran, S. (2011), Water quality studies through GIS at Bhavani Taulk Erode District, Tamilnadu, India. Eco. Env. & Cons. 17(2): 291-295.
- Prescott, G. M. (1982). Algae of the Western Great Lakes area. Otto Koeltz Scinces Publishers, West Germany.
- Purushothama, R., Mahesh Anand Goudar and Sayeswara, H.A. (20110). Seasonal Phytoplankton diversity and density in two lentic water bodies of Sagara, Karnataka, India. International Journal of Chemical Sciences (9(3); 1373-1390.
- Purushothama, R., Sayeswara, H.A. and Mahesh Anand Goudar (2011). Dynamics of Zooplankton diversity in relation to water quality of Heggere tank, Kanale, Sagar, Karnataka, India. Environment Conservation Jouranl. 12(1&2):29-34.
- Santhokumar Singh, A. Dakua, D. and Biswas, S. P. (2009). Physico-chemical parameters and fish enumeration of Maijan Beel (Wetland) of Upper Assam, Geobios, 36: 184-188.
- Sayeswara, H.A., Naik, K.L., Nafeesa Bugum and Ashashree, H.M. (2011). Potability of water inrelation ot some Physico-chemical parameter of Mudugodu pond, Chikkamagalur, Karnataka, India. Environment & Ecology. 29(1): 140-143.
- Sayeswara, H.A., Ravikumar Patil, H.S. and Mahesh Anand Goudar (2010). Studies on Physico-chemical parameters of Purle pond water of Shivamogga, Karnataka, India. International Journal Chemical Sciences 8(1): 582-588.
- Sayeswara, H.A., Vasantha Naik, T., Ravikumar Patil, H.S. and Mahesh Anand Goudar. (2012). Study on Habitat ecology and Phytoplankton diversity of Nagathibelagalu tank in Industrial town Bhadravathi, Karnataka, India. Nature Environment and Pollution Technology. 11(3): 419-423.
- Sayeswari, H.A., Mahesh Anand Goudar and Nafeesa Begum (2012a). Physico-chemical profile and Phytoplankton diversity of Murughamatta tank, Shivamogga, Karnataka, India. Pollution Research 31(3): 351-356.
- Technology and Arts Research Journal 2(2): 70-74, Trivedi, R.K., Goel, P. K. and Trishal, L. (1995). Practical methods in Ecology and Environmental Science,
- Vasantha Naik, 1 T., Puttaiah, E.T., Ravikumar Patil, H.S., Sayeswara, H.A. and Nafeesa Begum (2013). Phytoplankton community composition in Gowrikere tank, Anandapura, Sagara, Shivamogga, Karnataka, India. Science,